


National Marine Sanctuaries

Ocean Acidification: The Other CO₂ Problem

Richard A. Feely

NOAA/Pacific Marine Environmental Laboratory
February 2009

With special thanks to: James Orr, Victoria Fabry, Carol Turley, Chris Sabine, Joanie Kleypas, Kitack Lee, and Simone Alin






What we know about the ocean chemistry of ...*saturation state*



Saturation State

$$\Omega_{phase} = \frac{[Ca^{2+}][CO_3^{2-}]}{K_{sp,phase}^*}$$



$\Omega > 1 = \text{precipitation}$


$\Omega = 1 = \text{equilibrium}$

$\Omega < 1 = \text{dissolution}$





What we know about ocean CO₂ chemistry ...*from field observations*



WOCE/JGOFS/OACES Global CO₂ Survey
~72,000 sample locations
collected in the 1990s


DIC $\pm 2 \mu\text{mol kg}^{-1}$

TA $\pm 4 \mu\text{mol kg}^{-1}$

Sabine et al (2004)




What we know about ocean CO₂ chemistry...*from observed shoaling saturation horizons*





What we know about ocean CO₂ chemistry...*from observed aragonite and calcite saturation depths*




Feely et al. (2004)



Natural processes that could accelerate the ocean acidification of coastal waters



➤ Coastal Upwelling





North American Carbon Program

Continental Carbon Budgets, Dynamics, Processes, and Management





Aragonite Saturation State in west coast waters



Experiments on Many Scales

Biosphere 2



Provided by Mark Eakin



Aquaria and Small Mesocosms



SHARQ
Submersible Habitat for
Analyzing Reef Quality






Coccolithophore (single-celled algae)

$p\text{CO}_2$ 280-380 ppmv



Emiliana huxleyi

$p\text{CO}_2$ 780-850 ppmv




Calcification
decreased

- 9 to 18%



Gephyrocapsa oceanica



- 45%

Malformed liths at high CO_2

Manipulation of CO_2 system by addition of HCl or NaOH

Riebesell et al.(2000); Zondervan et al.(2001)



Shelled Pteropods (planktonic snails)


Respiratory CO_2 forced $\Omega_A < 1$

Shells of live animals start to dissolve within 48 hours

Whole shell:
Clio pyramidata

Arag. rods exposed

Prismatic layer
(1 μm) peels back




Aperture (~7 μm):
advanced dissolution

Normal shell: no
dissolution



Orr et al. (2005)

Response of mussels & oysters to elevated CO_2



Decrease in calcification rates for the 2 species:

Mytilus edulis

Crassostrea gigas

- Significant with pCO_2 increase and $[\text{CO}_3^{2-}]$ decrease


At pCO_2 740 ppmv:

- 25% decrease in calcification for mussels

- 10% decrease in calcification for oysters



Bivalve juvenile stages can also be sensitive to carbonate chemistry



Hard shell clam *Mercenaria*


- Common in soft bottom habitats

Used newly settled clams

- Size 0.3 mm
- Massive dissolution within 24 hours in undersaturated water; shell gone within 2 weeks
- Dissolution is source of mortality in estuaries & coastal habitats



Potential Effects on Open Ocean Food Webs



Coccolithophores



ARCOD@ims.uaf.edu

Copepods



Vicki Fabry

Pteropods



Barrie Kovish

Pacific Salmon




What we know about the biological impacts of ocean acidification *...on marine fish*

Research on Impacts of OA on Pacific Salmon

- ▲ Western Alaskan Sockeye
- ◆ British Columbia Sockeye
- Central Alaskan Pink
- Japanese Chum

Pink salmon diet















Predicted effect of climate change on pink salmon growth:

- 10% increase in water temperature leads to 3% drop in mature salmon body weight (physiological effect).
- 10% decrease in pteropod production leads to 20% drop in mature salmon body weight (prey limitation).



Scorecard of Biological Impacts

Physiological process	Major group	# species studied	Response to increasing CO ₂			
Calcification						
	Coccolithophores	4	2	1	1	1
	Planktonic Foraminifera	2	2	-	-	-
	Molluscs	4	4	-	-	-
	Echinoderms	2	2	-	-	-
	Tropical Corals	11	11	-	-	-
	Coralline Red Algae	1	1	-	-	-
Photosynthesis¹						
	Coccolithophores ²	2	-	2	2	-
	Prokaryotes	2	-	1	1	-
	Seagrasses	5	-	5	-	-
Nitrogen Fixation						
	Cyanobacteria	1	-	1	-	-
Reproduction						
	Molluscs	4	4	-	-	-
	Echinoderms	1	1	-	-	-

1) Strong interactive effects with nutrient and trace metals availability, light, and temperature
 2) Under nutrient replete conditions

Figure from Doney et al. (2009)



Conclusions

- Impacts of ocean acidification on ecosystems are largely unknown.
- Calcification in many planktonic organisms is reduced at elevated CO_2 , but the response is not uniform.
- Possible responses of ecosystems are speculative but could involve changes in species composition & abundances - could affect food webs, biogeochemical cycles.
- Baseline data with sufficient resolution are lacking in coastal regions where CaCO_3 saturation states are expected to decrease dramatically over in next 50-100 years.