

Received: 12 December 2018


Accepted: 7 May 2019


Published online: 06 June 2019

OPEN The vertical distribution and biological transport of marine microplastics across the epipelagic and mesopelagic water column

C. Anela Choyon, Bruce H. Robison, Tyler O. Gagneo, Benjamin Erwin, Evan Firl, Rolf U. Halden³, J. Andrew Hamilton¹, Kakani Katija¹, Susan E. Lisin², Charles Rolsky³ & Kyle S. Van Houtan 602,4

Plastic waste has been documented in nearly all types of marine environments and has been found in species spanning all levels of marine food webs. Within these marine environments, deep pelagic waters encompass the largest ecosystems on Earth. We lack a comprehensive understanding of the concentrations, cycling, and fate of plastic waste in sub-surface waters, constraining our ability to implement effective, large-scale policy and conservation strategies. We used remotely operated vehicles and engineered purpose-built samplers to collect and examine the distribution of microplastics in the Monterey Bay pelagic ecosystem at water column depths ranging from 5 to 1000 m. Laser Raman spectroscopy was used to identify microplastic particles collected from throughout the deep pelagic water column, with the highest concentrations present at depths between 200 and 600 m. Examination







The author of THE SEA AROUND US and
THE EDGE OF THE SEA
questions our attempt to control the
natural world about us

SILENT SPRING Rachel Carson



